

*The Physics  
of Time Travel*

*In memory of my  
father, who lives*

*Author: Javier Sánchez Sánchez*

# THE PHYSICS OF TIME TRAVEL

F. Javier Sánchez

*e-mail: javisan012@gmail.com*

## Abstract

The principle of General Covariance allows to construct the metric, the affine connection and the equations of the dynamics in accelerated reference frames, in the absence of gravity. The equations obtained explain the differences between gravity and inertia, the meaning of time and how to travel in time. After, the relativistic equations of motion of the rotating perfect fluid without pressure are solved, calculating the external force applied on it to go back in time. In the Appendix the formalism is generalized to the case in which gravity is present.

## INTRODUCTION

There are basically three ways in which time travel arises in General Relativity: rotating space-times containing Closed Timelike Curves [1-5], other geometries with CTCs [6-11] and space-times which allow for faster than light travel to generate CTCs [9-10], [12-14]. These models present several practical difficulties depending on the particular CTC containing geometry: the Tomimatsu-Sato geometries in vacuum involve a naked singularity, the CTC region in the Kerr space-time is confined behind an event horizon, superluminal travel often require substantial quantities of exotic matter [15], [16] and other models need infinitely large distributions of matter.

However, there is an alternative based on the Principle of Equivalence of Gravitation and Inertia postulated by Einstein in the General Relativity. As we know, it is easier to handle inertia appearing with the accelerated motion than gravity, although the effects in both cases must be locally equivalent. Due to this, it is possible to find metrics originated as a consequence of accelerated motion to travel to the past.

Unfortunately, there are different ways to deal with accelerated motion in relativity [17-27] and there is no general agreement on how to write the metric tensor of the accelerated reference frame not even the coordinate transformations between the inertial and accelerated frames. There is no agreement either on how to establish the relativistic transformation to rotating frames. The paper of Klauber reviews different theories of relativistic rotation [28].

We will follow a very common technique that consists of using the coordinate transformations between the inertial and rotating frames to construct the metric in the latter [28-34] with the same spatial coordinate transformations. The time coordinate transformation will be determined by the gauge. The gauge will be chosen to achieve time dilation, which guarantees validity to relativistic rotational velocities. As a result,

the time coordinate transformation depends on the only available degree of freedom, that is, the angular velocity of the rotation. After, the relativistic equations of motion of the fluid are easily obtained from the coordinate transformations or vice versa. Due to this, the validity of the coordinate transformations can be verified as exact solutions of the relativistic equations of motion. The principle of General Covariance provides physical meaning to this method.

This paper shows how the travel to the past is possible using a simple model of rotating perfect fluid without pressure. Firstly, it is assumed that the Gravitational field created by the fluid is negligible, which implies flat space-time. The time coordinate transformation between the inertial and rotating frames will be depend on the only available degree of freedom, that is, the angular velocity of the fluid, so this can be chosen so that the fluid goes back in time. The relativistic equations of motion of the fluid are easily obtained from the coordinate transformations checking that the external force applied on the fluid is always finite.

The main advantage is that in this flat space-time CTCs are not need to go back in time. Instead of that, the angular velocity of the fluid is chosen so that the time coordinate transformation between the inertial and rotating frames is a pointy function which allows the fluid to go back at any time in the past. The shape of this function is because travel to the past in a flat space-time must be understood as follow: on the rotating and inertial frames, time always goes forward. However, when the clocks on the rotating frame are compared with clocks on the inertial frame, the latter goes back at a rate determined by time dilation. In order to achieve this time reversal, the function must go from increasing to decreasing so that it must reach a pointy maximum. In addition, the travel to the past is possible because at time reversal, the fluid on the rotating reference frame can continue travelling to the future or it can travel to the past; but since the external force applied is finite and different in each case, an observer on the rotating frame can choose to travel to the past. Moreover, neither huge amounts of matter that curve space-time nor exotic matter or negative energy are needed and also the mass-energy is conserved in the Lorentzian frame of the observer at rest.

Finally we consider the case when the Gravitational field created by the fluid is important and therefore gravity is present.

Sections 2 to 5 consider the motion in a flat space-time without gravity. Section 2 studies the moving reference frames and the inertial field. Section 3, analyses the rotating reference frames. In section 4, the previous results are applied to obtain the relativistic motion of the rotating perfect fluid without pressure. In section 5, the force applied on this model of rotating fluid is calculated to go back in time. Finally, the Appendix connects with the equations of the field and gravity.

## I. MOVING REFERENCE FRAMES

There is no agreement on how to write the coordinate transformations between the inertial and accelerated frames. For this reason, the coordinate transformations used here to construct the metric are presented in a general form which validity will be determined as solutions of the relativistic equations of motion. Moreover, we will use a different gauge to constrain the available degrees of freedom and to achieve time dilation.

### Equations of Motion

When relativistic dynamics is studied, it is convenient to define two observers linked to two different reference frames: one that is considered at rest and the other moving with respect to the first. From now on they are going to be known as reference frame at rest  $S'$  and moving reference frame  $S$ .

Let  $S'$  be an inertial reference frame at rest in which the laws of Special Relativity are valid globally or at least within a finite volume. In this frame the metric is the Minkowski metric<sup>1</sup>

$$\eta_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \eta_{\nu\mu}. \quad (1)$$

An observer located in  $S'$  defines Cartesian coordinates  $x'^{\mu} = (t', x', y', z')$  to measure events. Let  $S$  be a moving reference frame with respect to  $S'$  in which other coordinates  $x^{\mu} = (x^0, x^1, x^2, x^3)$  are used to measure the events.

The equations of motion in  $S'$  of a fluid moving under the action of an external force are according to the laws of Special Relativity [35]

$$f'^{\mu} = \frac{\partial T'^{\mu\alpha}}{\partial x'^{\alpha}} \quad (2)$$

The equations of motion must be invariant under any coordinate transformations. Suppose that the coordinate transformations between  $S'$  and  $S$  are continuous functions of the form  $x'^{\mu} = x'^{\mu}(x^0, x^1, x^2, x^3)$ . These transformations must be global, covering all space-time or at least the volume of the fluid.

The inverse transformations are functions of the form  $x^{\mu} = x^{\mu}(t', x', y', z')$ . The necessary and sufficient condition for this to be true is that the Jacobian of the transformation must be nonzero.<sup>2</sup>

Then, carrying out any coordinate transformations from  $S'$  to  $S$ , in the equations of motion (2), we can find the equations of motion of a fluid in any reference frame, for example  $S$

<sup>1</sup> From now on, a system of units in which the speed of light is unity ( $c=1$ ) is used and any index that appears twice, once as a subscript and once as a superscript is understood to be summed over.

<sup>2</sup> Except maybe for some particular choices of coordinates which are called "singular".

$$f^\mu = T^{\mu\alpha}{}_{;\alpha} = \frac{\partial T^{\mu\alpha}}{\partial x^\alpha} + \Gamma_{\alpha\beta}^\alpha T^{\mu\beta} + \Gamma_{\alpha\beta}^\mu T^{\alpha\beta} \quad (3)$$

where

$$f^\mu = \frac{\partial x^\mu}{\partial x'^\alpha} f'^\alpha. \quad (4)$$

The metric is

$$g_{\mu\nu} = \eta_{\alpha\beta} \frac{\partial x'^\alpha}{\partial x^\mu} \frac{\partial x'^\beta}{\partial x^\nu} \quad (5)$$

and the affine connection is

$$\Gamma_{\mu\nu}^\lambda = \frac{\partial x^\lambda}{\partial x'^\alpha} \frac{\partial^2 x'^\alpha}{\partial x^\mu \partial x^\nu} \quad (6)$$

or

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\alpha} \left( \frac{\partial g_{\mu\alpha}}{\partial x^\nu} + \frac{\partial g_{\nu\alpha}}{\partial x^\mu} - \frac{\partial g_{\mu\nu}}{\partial x^\alpha} \right). \quad (7)$$

The principle of General Covariance says that equations of motion (3) are true in any reference frame in the absence of gravity. In practice, this happens when the gravitational field created by the fluid is so weak that it barely curves space-time. Nevertheless, they are the same as in a Gravitational field in the presence of external forces,<sup>35</sup> according to the Principle of Equivalence. It is important to note that they explicitly show the coordinate transformations between S' and S through the affine connection (6).

In order to solve the equations of motion (3) we need to fix the gauge, that is, the reference frame. If the reference frame S is chosen so that the fluid remains at rest with respect to it, then

$$\frac{dx^i}{d\tau} = 0. \quad (8)$$

Equations (8) do not fix the gauge alone. It is necessary to add another condition: the time-time component of the metric has to be invariant

$$g_{tt} = \eta_{tt} = -1. \quad (9)$$

This condition is needed for both the fluid and the reference frame S comove in time.

### Meaning of Time

As it is well known, the three spatial dimensions are characterized by freedom of motion. This is not so in the case of time as it is demonstrated below.

Equation (9) and the time-time component of Eq. (5) give

$$\left( \frac{\partial x'}{\partial t} \right)^2 + \left( \frac{\partial y'}{\partial t} \right)^2 + \left( \frac{\partial z'}{\partial t} \right)^2 = -1 + \left( \frac{\partial t'}{\partial t} \right)^2. \quad (10)$$

On the other hand, the components of the velocity of S with respect to S' are

$$v'^i = \frac{\frac{\partial x'^i}{\partial t}}{\frac{\partial t}{\partial t}} \quad (11)$$

since S is at rest in its own reference frame.

In addition, the magnitude of the velocity is calculated with Eqs. (11) and (10)

$$v' = \sqrt{v'^i v'^i} = \sqrt{\frac{-1 + \left(\frac{\partial t'}{\partial t}\right)^2}{\left(\frac{\partial t'}{\partial t}\right)^2}} \leq 1. \quad (12)$$

This equation shows, as expected, that it can never be greater than unity, that is, any reference frame can't move at a speed greater than the speed of light, with respect to S'. From Eq. (12), it is found

$$\frac{\partial t'}{\partial t} = \frac{1}{\sqrt{1 - v'^2}} = \gamma'. \quad (13)$$

This equation explains time dilation when velocity increases which guarantees validity to relativistic velocities and this is an experimental test to check the validity of Eq. (9). It shows that, in the absence of gravity, a clock at rest relative to another will run at the same rate, while one that moves relative to first will indicate its own time, different from the other. In the same way, motion in time depends on motion in space.<sup>3</sup> In

particular, the velocity of motion in time  $\frac{\partial t'}{\partial t}$  of S with respect to S' depends on its spatial velocity  $v'$ . In brief, time is a dimension in which there is no freedom of motion because motion in time is not independent of motion in space, but depends on it as in Eq. (13). This dependence makes it possible to determine the spatial velocity of a fluid to go back in time. The answer will be found in the following sections.

### Inertial Field

The curvature tensor in S can be defined as a function of the affine connection

$$R^\lambda{}_{\mu\nu\sigma} = \frac{\partial \Gamma^\lambda_{\mu\nu}}{\partial x^\sigma} - \frac{\partial \Gamma^\lambda_{\mu\sigma}}{\partial x^\nu} + \Gamma^\alpha_{\mu\nu} \Gamma^\lambda_{\sigma\alpha} - \Gamma^\alpha_{\mu\sigma} \Gamma^\lambda_{\nu\alpha}. \quad (14)$$

Since the curvature tensor in S' is clearly zero, it follows from the laws of transformation of the tensors that it is also zero in any other reference frame

$$R^\lambda{}_{\mu\nu\sigma} = 0. \quad (15)$$

This result implies that space-time in S is also flat instead of curved and that gravity does not appear either in the moving reference frame S.

The Ricci tensor is obtained by contraction of the curvature tensor and it is also zero according to Equation (15)

---

<sup>3</sup> As we know time also depends on the position in a gravitational field.

$$R_{\mu\nu} = R^{\alpha}{}_{\mu\alpha\nu} = 0. \quad (16)$$

This equation shows that the field in S has not sources but it is originated only by the motion of S with respect to S', that is, the inertial field has not sources which curve space-time.

## II. ROTATING REFERENCE FRAMES

There is no agreement either on how to establish the relativistic transformation to rotating frames. For this reason, the coordinate transformations used here to construct the metric in rotating frames are chosen using the only available degree of freedom and their validity will be determined as solutions of the relativistic equations of motion.

### Coordinate Transformations

A rotating reference frame S is an accelerated reference frame with an interesting property: it moves in the same place.

In this case it is convenient to define cylindrical coordinates  $x'' = (t, r, \theta, z)$  in S for practical reasons. If the motion is circular, it must add two equations

$$\left. \begin{aligned} x'^2 + y'^2 &= r^2 \\ z' &= z \end{aligned} \right\} \quad (17)$$

The coordinate transformations from the reference frame at rest S' to the rotating reference frame S compatible with Eqs. (17) are

$$\left. \begin{aligned} t' &= t'(t, r) \\ x' &= r \cos[\theta - \varphi(t)] \\ y' &= r \sin[\theta - \varphi(t)] \\ z' &= z \end{aligned} \right\} \quad (18)$$

The validity of these transformations can be verified as solutions of the relativistic equations of motion obtained in Section 4.1.

### The Metric Tensor

The metric in S is calculated with Eqs. (5) and (18)

$$\begin{aligned} g_{tt} &= -\left(\frac{\partial t'}{\partial t}\right)^2 + r^2 \left(\frac{d\varphi}{dt}\right)^2 = -1, g_{tr} = -\frac{\partial t'}{\partial t} \frac{\partial t'}{\partial r}, g_{t\theta} = -r^2 \frac{d\varphi}{dt}, g_{tz} = 0, \\ g_{rr} &= 1 - \left(\frac{\partial t'}{\partial r}\right)^2, g_{r\theta} = 0, g_{rz} = 0, \\ g_{\theta\theta} &= r^2, g_{\theta z} = 0, g_{zz} = 1 \end{aligned} \quad (19)$$

where Eq. (9) has been used in the first.

The first of Eqs. (19) can be written as

$$\frac{\partial t'}{\partial t} = \pm \sqrt{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}. \quad (20)$$

This equation has two solutions.<sup>4</sup> The positive solution implies that  $t'$  increases with  $t$ , which is known as time dilation. In section 5 we will see when negative solution is admissible, which ensures a decrease of  $t'$  as  $t$  increases, which allows to go back in time.

### The Affine Connection

The affine connection is calculated through Eqs. (7), with the inverse of the metric tensor given by

$$g^{\mu\nu} = \frac{\partial x^\mu}{\partial x'^\alpha} \frac{\partial x^\nu}{\partial x'^\beta} \eta^{\alpha\beta}, \quad (21)$$

the metric (19) and its derivatives

$$\begin{aligned} \Gamma_{tt}^r &= -r \left( \frac{d\varphi}{dt} \right)^2, \Gamma_{tt}^t = \frac{r \left( \frac{d\varphi}{dt} \right)^2}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \Gamma_{\theta t}^t = -\frac{r \frac{\partial t'}{\partial t} \frac{\partial t'}{\partial r} \frac{d\varphi}{dt}}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \\ \Gamma_{rr}^t &= \frac{\frac{\partial t'}{\partial t} \frac{\partial^2 t'}{\partial r^2}}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \Gamma_{\theta\theta}^t = \frac{r \frac{\partial t'}{\partial t} \frac{\partial t'}{\partial r}}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \Gamma_{t\theta}^r = r \frac{d\varphi}{dt}, \\ \Gamma_{tr}^\theta &= -\frac{\frac{d\varphi}{dt}}{r \left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)^2 \right]}, \Gamma_{t\theta}^\theta = -\frac{r \frac{\partial t'}{\partial t} \frac{\partial t'}{\partial r} \left( \frac{d\varphi}{dt} \right)^2}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \Gamma_{\theta\theta}^r = -r, \\ \Gamma_{rr}^\theta &= \frac{\frac{\partial t'}{\partial t} \frac{\partial^2 t'}{\partial r^2} \frac{d\varphi}{dt}}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \Gamma_{r\theta}^\theta = \frac{1}{r}, \Gamma_{\theta\theta}^\theta = \frac{r \frac{\partial t'}{\partial t} \frac{\partial t'}{\partial r} \frac{d\varphi}{dt}}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}. \end{aligned} \quad (22)$$

The rest of the components are zero.

## III. ROTATING PERFECT FLUID WITHOUT PRESSURE

### Equations of Motion in the Rotating Reference Frame

---

<sup>4</sup> From now on, both signs will be shown except in the case when the choice of one of them is justified.



In this section we apply the above results to the simplest model of fluid, that is, a perfect fluid without pressure. In this case the energy-momentum tensor of the perfect fluid is

$$T^{\mu\nu} = \rho \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}. \quad (23)$$

If the fluid is at rest in the rotating reference frame S, Eqs. (8), (9) and (23) give

$$T^{tt} = \rho; \quad T^{ti} = T^{it} = 0; \quad T^{ij} = 0. \quad (24)$$

In addition, if the mass-energy is conserved in S' the continuity equation of the fluid must be<sup>5</sup>

$$f'^{\mu} = 0 \quad (25)$$

and since the components of the density of the external force  $f'^{\mu}$  are calculated through Eqs. (4)

$$f'^{\mu} = \frac{\partial x'^{\mu}}{\partial x^{\alpha}} f^{\alpha} \quad (26)$$

the Eq. (25) give

$$\rho = \rho_i \frac{1 + r^2 \left( \frac{d\varphi}{dt} \right)_{t=t_i}^2}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}. \quad (27)$$

This equation shows that the proper energy density of the fluid  $\rho$  is always positive or zero.

The components of the density of the external force applied to the fluid in S are calculated using Eqs. (3), (22), (24) and (27)<sup>6</sup>

$$f^t = \pm \rho_i r \frac{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)_{t=t_i}^2 \right] \sqrt{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2} \frac{\partial t'}{\partial r} \left( \frac{d\varphi}{dt} \right)^2}{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)^2 \right]^2}, \quad (28)$$

$$f^r = -\rho_i r \frac{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)_{t=t_i}^2 \right] \left( \frac{d\varphi}{dt} \right)^2}{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}, \quad (29)$$

$$f^\theta = \rho_i \frac{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)_{t=t_i}^2 \right] \left[ \pm r \sqrt{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2} \frac{\partial t'}{\partial r} \left( \frac{d\varphi}{dt} \right)^3 - \frac{d^2\varphi}{dt^2} \right]}{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)^2 \right]^2}, \quad (30)$$

<sup>5</sup> This equation is not covariant so it is only valid in the reference frame at rest S'.

<sup>6</sup> The density of force in the S.I. is obtained dividing by  $c^2$  taking  $c \approx 3 \cdot 10^8$  m/s.

$$f^z = 0. \quad (31)$$

The rotational motion is only due to the radial  $f^r$  and the tangential  $f^\theta$  components of the density of the external force given by Eqs. (29) and (30) and from now on, only these two components will be referred to.

Introducing Eqs. (28) to (31) into Eqs. (3), we check that the coordinate transformations Eqs. (18) are solutions of the relativistic equations of motion (3) in the gauge given by Eqs. (8) and (9) with the constraints Eqs. (17), (25) and the equation of the invariant interval.

### Equations of Motion in the Reference Frame at Rest

It is also convenient to determine the motion of the rotating fluid in the reference frame  $S'$ . The inverses of Eqs. (18) allow to obtain the density of the external force directly in  $S'$  using Eqs. (25), (26), and (28) to (31)

$$f'^t = 0, \quad (32)$$

$$f'^r = f^r, \quad (33)$$

$$f'^\theta = -\rho_i \frac{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)_{t=t_i}^2 \right] \frac{d^2\varphi}{dt^2}}{\left[ 1 + r^2 \left( \frac{d\varphi}{dt} \right)^2 \right]^2}, \quad (34)$$

$$f'^z = f^z = 0. \quad (35)$$

Also for the same reasons given for  $S$ , from now on only the radial  $f'^r$  and tangential  $f'^\theta$  components of the density of the external force will be referred to.

In addition, introducing Eq. (20) into Eq. (12) gives the velocity of  $S$

$$v' = \frac{r \frac{d\varphi}{dt}}{\sqrt{1 + r^2 \left( \frac{d\varphi}{dt} \right)^2}} \leq 1. \quad (36)$$

If the angular velocity is constant  $\frac{d\varphi}{dt} = \omega$  and the fluid is initially at rest at  $t_i = t_0$  in  $S$  in which the density is  $\rho_0$ , then the relativistic centripetal force given by Eqs. (33) and (29)<sup>7</sup>

$$f'_{rel}{}^r = -\rho_0 \frac{\omega^2 r}{1 + \omega^2 r^2} \quad (37)$$

<sup>7</sup>  $\rho_0$  can be chosen as the proper energy density of water, that is  $\rho_0 = 27 \cdot 10^{27} \text{ Kg} / \text{s}^3 \approx 9 \cdot 10^{19} \text{ J} / \text{m}^3$ .




Fig. 1.  $f''_{rel}$  as function of  $r$  with constant angular velocity  $\omega = 3 \text{ rad/s}$ . The minimum is  $f''_{rel} = -4.05 \cdot 10^{28} \text{ Kg/s}^4 = -4.5 \cdot 10^{11} \text{ N/m}^3$  at  $r = 1/\omega$ .

is different from the classical law

$$f'_{clas}{}^r = -\rho_0 \omega^2 r. \quad (38)$$




Fig. 2. Classical law for  $\omega = 3 \text{ rad/s}$ .  $f''_{clas} = -8.1 \cdot 10^{28} \text{ kg/s}^4 = -9 \cdot 10^{11} \text{ N/m}^3$  at  $r = 1/\omega$ .

#### IV. TIME TRAVEL

In this section we apply the results obtained in the previous sections to determine the dynamics of the rotating perfect fluid without pressure to go back in time. This means finding the external force applied on the fluid to go back in time with rotational motion. For this purpose the only available degree of freedom of a rotating perfect fluid, that is, the angular velocity  $\frac{d\varphi}{dt}$  is fixed. This variable is chosen so that the associated time shift obtained from Eq. (20) allows the fluid to go back in time. The travel will take five

legs. The procedure is the following: first, we fix the angular velocity  $\frac{d\varphi}{dt}$  and second, we calculate  $\frac{\partial t'}{\partial t}$  through Eq. (20), integrating the last we obtain  $t' = t'(t, r)$  and deriving this we calculate  $\frac{\partial t'}{\partial r}$ . In addition, we use Eq. (27). Then, we insert these results into the equations of motion (29) and (30). Occasionally, we will use Eqs. (33) and (34) to show the motion in  $S'$ .

### Temporal Acceleration

At the beginning, the observers located in  $S'$  and  $S$  are in an inertial reference frame in which the Minkowski metric is valid. This allows to synchronise the clocks of both frames according to the criterion of the light signals proposed by Einstein [36].

The first leg of time travel occurs during the interval  $t_0 \leq t \leq t_1$  for an observer located in  $S$ . If the reference frame  $S$  is initially at rest at  $t = t_0$  the simplest function  $\frac{d\varphi}{dt}$  is

$$\frac{d\varphi_I}{dt} = a_1(t - t_0) \quad (39)$$

where  $a_1$  is a constant that can be chosen so that

$$a_1 > 0. \quad (40)$$

In addition

$$\frac{\partial t'_I}{\partial t} = \sqrt{1 + a_1^2 r^2 (t - t_0)^2}, \quad (41)$$

$$t'_I = t'_0 + \frac{t - t_0}{2} \sqrt{1 + a_1^2 r^2 (t - t_0)^2} + \frac{1}{2a_1 r} \text{Ln} \left[ a_1 r (t - t_0) + \sqrt{1 + a_1^2 r^2 (t - t_0)^2} \right] \quad (42)$$

and

$$\frac{\partial t'_I}{\partial r} = \frac{t - t_0}{2r} \sqrt{1 + a_1^2 r^2 (t - t_0)^2} - \frac{1}{2a_1 r^2} \text{Ln} \left[ a_1 r (t - t_0) + \sqrt{1 + a_1^2 r^2 (t - t_0)^2} \right] \geq 0. \quad (43)$$

We obtain for this fluid in the reference frame  $S$  during the first leg of the time travel

$$f_I^r = - \frac{\rho_0 a_1^2 r (t - t_0)^2}{1 + a_1^2 r^2 (t - t_0)^2}, \quad (44)$$

$$f_I^\theta = \rho_0 \frac{r \sqrt{1 + a_1^2 r^2 (t - t_0)^2} \frac{\partial t'_I}{\partial r} a_1^3 (t - t_0)^3 - a_1}{[1 + a_1^2 r^2 (t - t_0)^2]^2} \quad (45)$$

where  $\frac{\partial t'_I}{\partial r}$  is given by Eq. (43).

### Time Reversal

The second leg of time travel occurs during the interval  $t_1 \leq t \leq t_3$  for an observer located in S. The simplest function  $\frac{d\varphi}{dt}$  which gives time reversal is

$$\frac{d\varphi_{II}}{dt} = \frac{1}{(t_2 - t)^{\frac{1}{3}}} \sqrt{a_2(t_2 - t)^{\frac{2}{3}} + b_2}. \quad (46)$$

Introducing Eq. (46) into Eq. (20) we have

$$\frac{\partial t'_{II}}{\partial t} = \frac{1}{(t_2 - t)^{\frac{1}{3}}} \sqrt{(1 + a_2 r^2)(t_2 - t)^{\frac{2}{3}} + b_2 r^2}. \quad (47)$$

It is observed that  $\frac{\partial t'_{II}}{\partial t}$  changes sign at  $t = t_2$ , which allows to go back in time as it was discussed in Eq. (20). Integrating Eq. (47) yields

$$t'_{II} = t'_1 + \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2)(t_2 - t_1)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} - \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2)(t_2 - t)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} \quad (48)$$

where the function  $t'_1 = t'_1(r)$  is calculated from the continuity of  $t'$

$$t'_1 = t'_0 + \frac{t_1 - t_0}{2} \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} + \frac{1}{2a_1 r} \text{Ln} \left[ a_1 r (t_1 - t_0) + \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} \right]. \quad (49)$$

In order to avoid divergences it is convenient that

$$a_2 > 0. \quad (50)$$

The shape of the function Eq. (48) that reaches a maximum at  $t = t_2$  allows time reversal which occurs at that time.




Fig 3.  $t'_{II}$  as function of  $t$  taking  $r = r_0 = 10^{-7} s = 30m$  during the second leg of time travel. The constants can be chosen as:  $a_1 = 1 \text{ rad/s}^2$ ,  $a_2 = 10^{14} \text{ rad}^2 / \text{s}^2$ ,  $b_2 = 10^{14} \text{ rad}^2 / \text{s}^{4/3}$ ,  $t_0 = 0$ ,  $t_1 = 1s$ ,  $t_2 = 2s$  and  $t'_0 = 0$ .

Deriving Eq. (48) and taking into account Eq. (49) we obtain

$$\frac{\partial t'_{II}}{\partial r} = \frac{t_1 - t_0}{2r} \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} - \frac{1}{2a_1 r^2} \text{Ln} \left[ a_1 r (t_1 - t_0) + \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} \right]$$

$$\begin{aligned}
 & + \frac{a_2 r (1 + a_2 r^2) (t_2 - t_1)^{\frac{2}{3}} + a_2 b_2 r^3 + 3b_2 r \sqrt{(1 + a_2 r^2) (t_2 - t_1)^{\frac{2}{3}} + b_2 r^2}}{(1 + a_2 r^2)^2} \\
 & - \frac{a_2 r (1 + a_2 r^2) (t_2 - t)^{\frac{2}{3}} + a_2 b_2 r^3 + 3b_2 r \sqrt{(1 + a_2 r^2) (t_2 - t)^{\frac{2}{3}} + b_2 r^2}}{(1 + a_2 r^2)^2} > 0.
 \end{aligned} \tag{51}$$

The proper energy density at time  $t = t_1$  is obtained from Eq. (27)

$$\rho_1 = \rho(t_1) = \frac{\rho_0}{1 + r^2 \left( \frac{d\varphi_I}{dt} \right)_{t=t_1}^2}. \tag{52}$$


Introducing Eqs. (46) and (52) into Eqs. (29) and (30) we obtain for this fluid in the reference frame S during the second leg of the time travel

$$f_{II}^r = -\rho_0 r \frac{a_2 (t_2 - t)^{\frac{2}{3}} + b_2}{(1 + a_2 r^2) (t_2 - t)^{\frac{2}{3}} + b_2 r^2}, \tag{53}$$

$$f_{II}^\theta = \rho_0 \frac{r \sqrt{(1 + a_2 r^2) (t_2 - t)^{\frac{2}{3}} + b_2 r^2} \frac{\partial t'_{II}}{\partial r} \left[ a_2 (t_2 - t)^{\frac{2}{3}} + b_2 \right]^{\frac{3}{2}} - \frac{b_2}{3 \sqrt{a_2 (t_2 - t)^{\frac{2}{3}} + b_2}}}{\left[ (1 + a_2 r^2) (t_2 - t)^{\frac{2}{3}} + b_2 r^2 \right]^2} \tag{54}$$

where  $\frac{\partial t'_{II}}{\partial r}$  is given by Eq. (51). It is observed that  $f_{II}^r$  and  $f_{II}^\theta$  are finite through the interval although the metric in Eq. (19) is singular at  $t = t_2$  in spite of the metric in S is the Minkowski metric, free of singularities.

a)



b)




Fig 4. If we take  $r = r_0 = 10^{-7} s = 30m$ ,  $a_2 = 10^{14} rad^2 / s^2$ ,  $b_2 = 10^{14} rad^2 / s^{4/3}$  and  $\rho_0 = 27 \cdot 10^{27} Kg / s^3 \approx 9 \cdot 10^{19} J / m^3$  we obtain: a)  $f_{II}^r$  as function of  $t$  during the second leg of time travel. The minimum is  $f_{II}^r = -2.7 \cdot 10^{35} kg / s^4 = -3 \cdot 10^{18} N / m^3$  at  $t = t_2 = 2s$ . b)  $f_{II}^\theta$  as function of  $t$  during the second leg of time travel. The maximum is  $f_{II}^\theta = 4.315 \cdot 10^{49} kg / s^4 = 4.794 \cdot 10^{32} N / m^3$  at  $t = t_2 = 2s$ .

In order to obtain Eq. (54) when  $t \geq t_2$ , the negative sign in Eq. (20) has been chosen to allow the travel to the past. This choice is possible because the density of the external force  $f_{II}^\theta$  in Eq. (30) or (54) depends on the sign of  $\frac{\partial t'_{II}}{\partial t}$  since  $\frac{\partial t'_{II}}{\partial r} > 0$ . This means that in the reference frame S, different values of  $f_{II}^\theta$  are obtained according to the positive or negative sign of  $\frac{\partial t'_{II}}{\partial t}$ . This makes the trip to the past possible, because when  $t \geq t_2$ , the positive and negative solutions in Eq. (20) are physically different in S, which allows to choose between one of them.

It can also be verified that changing the sign of the solution of Eq. (54) also allows the trip to the past when  $t \geq t_2$ . Both solutions only differ in the sign of the angular velocity  $\frac{d\varphi_{II}}{dt}$ .

The second leg of time travel occurs during the interval  $t'_1 \leq t' \leq t'_3$  for an observer located in S'.

The rotating motion of this fluid in this reference frame during this leg is

$$f_{II}^{r'} = f_{II}^r = -\rho_0 r \frac{a_2(t_2 - t)^{\frac{2}{3}} + b_2}{(1 + a_2 r^2)(t_2 - t)^{\frac{2}{3}} + b_2 r^2}, \quad (55)$$

$$f_{II}^{\theta'} = -\frac{\rho_0 b_2}{3\sqrt{a_2(t_2 - t)^{\frac{2}{3}} + b_2} \left[ (1 + a_2 r^2)(t_2 - t)^{\frac{2}{3}} + b_2 r^2 \right]^2}. \quad (56)$$




Fig. 5.  $f_{II}^{\theta'}$  as function of  $t$  taking  $r = r_0 = 10^{-7} s = 30m$ ,  $a_2 = 10^{14} rad^2 / s^2$ ,  $b_2 = 10^{14} rad^2 / s^{4/3}$  and  $\rho_0 = 27 \cdot 10^{27} Kg / s^3 \approx 9 \cdot 10^{19} J / m^3$  during the second leg of time travel. The minimum is  $f_{II}^{\theta'} = -9 \cdot 10^{34} kg / s^4 = -10^{18} N / m^3$  at  $t = t_2 = 2s$ .

We also note that  $f_{II}^{r'}$  and  $f_{II}^{\theta'}$  are finite through the interval.

It is observed that the density of the external force  $f_{II}^{\prime\theta}$  given by Eq. (34) or (56) does not depend on  $\frac{\partial t'_{II}}{\partial t}$ . This means that an observer located in  $S'$  can not force the time backwards because the positive and negative solutions of Eq. (20) lead to the same  $f_{II}^{\prime\theta}$ . For this reason,  $f_{II}^{\prime\theta}$  only provides  $\frac{\partial t'_{II}}{\partial t} > 0$ . On the other hand, an observer located in the reference frame  $S$ , physically distinguishes both solutions and he can choose the solution in Eq. (30) or (54) that causes the time reversal. It can be advisable to write Eqs. (55) and (56) as a function of time  $t'$  measured in  $S'$ . For this, it is necessary to find  $t$  as a function of  $t'$  from Eq. (48). It can be checked by inspection of Eqs. (36), (46) and (27) that the fluid reaches the speed of light in  $S'$  at time  $t = t_2$  because the proper energy density of the fluid is null at that time in which time reversal occurs.

### Back to the Past

The third leg of time travel occurs during the interval  $t_3 \leq t \leq t_4$  for an observer located in  $S$ . It is convenient to keep angular velocity  $\frac{d\varphi}{dt}$  constant during this leg and thus

$$\frac{d\varphi_{III}}{dt} = a_3, \quad (57)$$

where  $a_3$  is a constant so that

$$a_3 < 0. \quad (58)$$

In addition

$$\frac{\partial t'_{III}}{\partial t} = \sqrt{1 + a_3^2 r^2} \quad (59)$$

and

$$t'_{III} = t'_3 + \sqrt{1 + a_3^2 r^2} (t - t_3) \quad (60)$$

where

$$t'_3 = t'_0 + \frac{t_1 - t_0}{2} \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} + \frac{1}{2a_1 r} \operatorname{Ln} \left[ a_1 r (t_1 - t_0) + \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} \right] + \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2) (t_2 - t_1)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} - \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2) (t_2 - t_3)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}}. \quad (61)$$

Deriving Eq. (60) we obtain



$$\frac{\partial t'_{III}}{\partial r} = \frac{dt'_3}{dr} + \frac{a_3^2 r}{\sqrt{1+a_3^2 r^2}}(t-t_3). \quad (62)$$

It is necessary to determine the velocity (59) of the travel backward in time. As  $\frac{\partial t'_{III}}{\partial t}$  depends on  $r$  to determine this velocity it is convenient to fix  $r$  so that, in  $r=r_0$  verifies

$$\left. \frac{\partial t'_{III}}{\partial t} \right|_{r_0, t_3} = -V \quad (63)$$

where  $V$  is a constant greater than zero

$$V > 0. \quad (64)$$

We obtain for this fluid in the reference frame S during the third leg of time travel

$$f_{III}^r = -\rho_0 r \frac{a_3^2}{1+a_3^2 r^2}, \quad (65)$$

$$f_{III}^\theta = \rho_0 \frac{r \sqrt{1+a_3^2 r^2} \frac{\partial t'_{III}}{\partial r} a_3^3}{[1+a_3^2 r^2]^2} \quad (66)$$

where  $\frac{\partial t'_{III}}{\partial r}$  is given by Eqs. (61) and (62).

### New Time Reversal

The fourth leg of time travel occurs during the interval  $t_4 \leq t \leq t_6$  for an observer located in S. The simplest function  $\frac{d\varphi}{dt}$  which gives a new time reversal is

$$\frac{d\varphi_{IV}}{dt} = \frac{1}{(t-t_5)^{\frac{1}{3}}} \sqrt{a_4(t-t_5)^{\frac{2}{3}} + b_4}. \quad (67)$$

where  $t_5$ ,  $a_4$  and  $b_4$  are constants verifying

$$t_4 < t_5 < t_6; \quad b_4 > 0; \quad b_4 > -a_4(t_4 - t_5)^{\frac{2}{3}}. \quad (68)$$

Introducing Eq. (67) into Eq. (20) we have

$$\frac{\partial t'_{IV}}{\partial t} = \frac{1}{(t-t_5)^{\frac{1}{3}}} \sqrt{(1+a_4 r^2)(t-t_5)^{\frac{2}{3}} + b_4 r^2}. \quad (69)$$

It is observed that  $\frac{\partial t'_{IV}}{\partial t}$  changes sign at  $t=t_5$  as it was discussed in Eq. (20).

Integrating Eq. (69) yields

$$t'_{IV} = t'_4 - \frac{1}{1+a_4 r^2} \left[ (1+a_4 r^2)(t-t_5)^{\frac{2}{3}} + b_4 r^2 \right]^{\frac{3}{2}} + \frac{1}{1+a_4 r^2} \left[ (1+a_4 r^2)(t-t_5)^{\frac{2}{3}} + b_4 r^2 \right]^{\frac{3}{2}} \quad (70)$$

where

$$\begin{aligned}
 t'_4 = t'_0 + \frac{t_1 - t_0}{2} \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} + \frac{1}{2a_1 r} \operatorname{Ln} \left[ a_1 r (t_1 - t_0) + \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} \right] \\
 - \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2) (t_2 - t_3)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} + \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2) (t_2 - t_1)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} \\
 + \frac{1}{(t_2 - t_3)^{\frac{1}{3}}} \sqrt{(1 + a_2 r^2) (t_2 - t_3)^{\frac{2}{3}} + b_2 r^2} (t_4 - t_3). \tag{71}
 \end{aligned}$$

In order to avoid divergences in Eq. (70) it is convenient that

$$a_4 > 0. \tag{72}$$

The shape of the function (70) that reaches a minimum at  $t = t_5$  allows a new time reversal which occurs at that time.

Deriving Eq. (70) we obtain

$$\begin{aligned}
 \frac{\partial t'_{IV}}{\partial r} = \frac{dt'_4}{dr} - \frac{a_4 r (1 + a_4 r^2) (t_4 - t_5)^{\frac{2}{3}} + a_4 b_4 r^3 + 3b_4 r}{(1 + a_4 r^2)^2} \sqrt{(1 + a_4 r^2) (t_4 - t_5)^{\frac{2}{3}} + b_4 r^2} \\
 + \frac{a_4 r (1 + a_4 r^2) (t - t_5)^{\frac{2}{3}} + a_4 b_4 r^3 + 3b_4 r}{(1 + a_4 r^2)^2} \sqrt{(1 + a_4 r^2) (t - t_5)^{\frac{2}{3}} + b_4 r^2} < 0. \tag{73}
 \end{aligned}$$

The proper energy density at time  $t = t_4$  is calculated from Eq. (27)

$$\rho_4 = \rho(t_4) = \frac{\rho_0}{1 + r^2 \left( \frac{d\varphi_{III}}{dt} \right)_{t=t_4}^2}. \tag{74}$$

Introducing Eqs. (67) and (74) into Eqs. (29) and (30) we obtain for this fluid in the reference frame S during the fourth leg of time travel

$$f_{IV}^r = -\rho_0 r \frac{a_4 (t - t_5)^{\frac{2}{3}} + b_4}{(1 + a_4 r^2) (t - t_5)^{\frac{2}{3}} + b_4 r^2}, \tag{75}$$

$$\begin{aligned}
 f_{IV}^\theta = \rho_0 \frac{r \sqrt{(1 + a_4 r^2) (t - t_5)^{\frac{2}{3}} + b_4 r^2} \frac{\partial t'_{IV}}{\partial r} \left[ a_4 (t - t_5)^{\frac{2}{3}} + b_4 \right]^{\frac{3}{2}} + \frac{b_4}{3 \sqrt{a_4 (t - t_5)^{\frac{2}{3}} + b_4}}}{\left[ (1 + a_4 r^2) (t - t_5)^{\frac{2}{3}} + b_4 r^2 \right]^2} \tag{76}
 \end{aligned}$$

where  $\frac{\partial t'_{IV}}{\partial r}$  is given by Eqs. (71) and (73). We also note that  $f_{IV}^r$  and  $f_{IV}^\theta$  are finite through the interval although the metric in Eq. (19) is singular at  $t = t_5$ .

Again, it is observed that the new time reversal is possible because  $f_{IV}^\theta$  depends on the

sign of  $\frac{\partial t'_{IV}}{\partial t}$  because  $\frac{\partial t'_{IV}}{\partial r} < 0$ .

We can prove that  $f''_{IV}$  and  $f'^{\theta}_{IV}$  are finite through the interval. Again, it can be verified that at time  $t = t_5$  in S, the fluid reaches the speed of light in S'.

### Temporal Deceleration

The fifth leg of time travel occurs during the interval  $t_6 \leq t \leq T$  for an observer located in S. If the rotation stops at the end of the travel, at time  $t = T$  in S, the simplest function  $\frac{d\varphi}{dt}$  is

$$\frac{d\varphi_V}{dt} = a_5(t - T) \quad (77)$$

where  $a_5$  is a constant such that

$$a_5 < 0. \quad (78)$$

In addition

$$\frac{\partial t'_V}{\partial t} = \sqrt{1 + a_5^2 r^2 (t - T)^2} \quad (79)$$

and

$$\begin{aligned} t'_V = t'_6 - \frac{t_6 - T}{2} \sqrt{1 + a_5^2 r^2 (t_6 - T)^2} - \frac{1}{2a_5 r} \text{Ln} \left[ a_5 r (t_6 - T) + \sqrt{1 + a_5^2 r^2 (t_6 - T)^2} \right] \\ + \frac{t - T}{2} \sqrt{1 + a_5^2 r^2 (t - T)^2} + \frac{1}{2a_5 r} \text{Ln} \left[ a_5 r (t - T) + \sqrt{1 + a_5^2 r^2 (t - T)^2} \right]. \end{aligned} \quad (80)$$

where

$$\begin{aligned} t'_6 = t'_0 + \frac{t_1 - t_0}{2} \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} + \frac{1}{2a_1 r} \text{Ln} \left[ a_1 r (t_1 - t_0) + \sqrt{1 + a_1^2 r^2 (t_1 - t_0)^2} \right] \\ - \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2)(t_2 - t_3)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} + \frac{1}{1 + a_2 r^2} \left[ (1 + a_2 r^2)(t_2 - t_1)^{\frac{2}{3}} + b_2 r^2 \right]^{\frac{3}{2}} \\ + \frac{1}{(t_2 - t_3)^{\frac{1}{3}}} \sqrt{(1 + a_2 r^2)(t_2 - t_3)^{\frac{2}{3}} + b_2 r^2} (t_4 - t_3) + \frac{1}{1 + a_4 r^2} \left[ (1 + a_4 r^2)(t_6 - t_5)^{\frac{2}{3}} + b_4 r^2 \right]^{\frac{3}{2}} \\ - \frac{1}{1 + a_4 r^2} \left[ (1 + a_4 r^2)(t_4 - t_5)^{\frac{2}{3}} + b_4 r^2 \right]^{\frac{3}{2}}. \end{aligned} \quad (81)$$

It is convenient that the travel ends at a certain time  $T'$  in the past of S'. As  $t' = t'(t, r)$ , it is advisable to fix  $r$  in which this happens. This will be  $r = r_0$  and so

$$t'_V(T)_{r_0} = T' \quad (82)$$

Deriving Eq. (80) we obtain

$$\frac{\partial t'_V}{\partial r} = \frac{dt'_6}{dr} - \frac{t_6 - T}{2r} \sqrt{1 + a_5^2 r^2 (t_6 - T)^2} + \frac{1}{2a_5 r^2} \text{Ln} \left[ a_5 r (t_6 - T) + \sqrt{1 + a_5^2 r^2 (t_6 - T)^2} \right]$$

$$+ \frac{t-T}{2r} \sqrt{1+a_5^2 r^2 (t-T)^2} - \frac{1}{2a_5 r^2} \text{Ln} \left[ a_5 r (t-T) + \sqrt{1+a_5^2 r^2 (t-T)^2} \right]. \quad (83)$$

We obtain for this fluid in the reference frame S during the fifth leg of time travel

$$f_V^r = -\frac{\rho_0 a_5^2 r (t-T)^2}{1+a_5^2 r^2 (t-T)^2}, \quad (84)$$

$$f_V^\theta = \rho_0 \frac{r \sqrt{1+a_5^2 r^2 (t-T)^2} \frac{\partial t_V'}{\partial r} a_5^3 (t-T)^3 - a_5}{[1+a_5^2 r^2 (t-T)^2]^2} \quad (85)$$

where  $\frac{\partial t_V'}{\partial r}$  is given by Eqs. (81) and (83). Observe that the constants appearing in the equations have been chosen to avoid divergences and they will be constrained by Eqs. (63), (82) and continuity conditions.

And this is the end of the travel to the past. As it is easy to imagine, the return travel to the present can be carried out in a similar way in only three legs that do not entail as much difficulty as the travel to the past.

Finally, note that the travel to the past can be used in interstellar space travel. Indeed, due to the effect of time dilation, if a traveller reached a speed close to that of light he could reach his destination star in a few months. However, when he returned to Earth he will note that here hundreds or thousands of years have passed. If the traveler went back in time he could make the duration of the trip the same for both the traveller and people on Earth.

## APPENDIX: EINSTEIN'S EQUATIONS WITH SCALAR FIELD

The above results are adequate in a flat space-time in the absence of gravity when the gravitational field created by the fluid is so weak that it barely curves space-time which happens in practice. Although it is not relevant in the exposed developments, it is convenient to generalize the formalism to the case in which the gravitational field generated by the fluid is important. The results in the absence of gravity are obtained as a particular case. To do this, it is necessary to resort to Einstein's field equations.

First, it should be noted that the energy-momentum tensor (23) is not conserved, as it is deduced from Eq. (3), so that it is not compatible with Einstein's equations

$$R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R = -8\pi G T^{\mu\nu} \quad (A.1)$$

for which the energy-momentum tensor is conserved. This means that the energy-momentum tensor (23) is not adequate to calculate the gravitational field created by a perfect fluid subjected to mechanical forces.

The simplest way to avoid this difficulty is introducing a scalar field  $\phi$  into Einstein's Eqs. (A.1) similarly to the Brans-Dicke Theory [37]. This scalar field verifies a new differential equation. The simplest generally covariant field equation for such a scalar field is

$$\phi_{;\alpha}^{\alpha} = 4\pi\lambda T_{M\alpha}^{\alpha} \quad (\text{A.2})$$

where  $\lambda$  is a coupling constant and  $T_M^{\mu\nu}$  is the energy-momentum tensor of the matter. The most general symmetric tensor for this field must contain terms which involves two derivatives of one or two  $\phi$  fields

$$T_{\phi}^{\mu\nu} = A(x)\phi_{;\mu}^{\mu}\phi_{;\nu}^{\nu} + B(x)g^{\mu\nu}\phi_{;\alpha}^{\alpha}\phi_{;\alpha}^{\alpha} + C(x)\phi_{;\mu}^{\mu;\nu} + D(x)g^{\mu\nu}\phi_{;\alpha}^{\alpha}. \quad (\text{A.3})$$

Einstein's Eqs. (A.1) including this scalar field are

$$R^{\mu\nu} - \frac{1}{2}g^{\mu\nu}R = -8\pi G(T_M^{\mu\nu} + T_{\phi}^{\mu\nu}) \quad (\text{A.4})$$

where  $T_M^{\mu\nu}$  verifies

$$f^{\mu} = T_M^{\mu\alpha}_{;\alpha} \quad (\text{A.5})$$

and  $T_{\phi}^{\mu\nu}$  is the energy-momentum tensor of the scalar field given by Eqs. (A.3).

In order to determine the functions  $A(x)$ ,  $B(x)$ ,  $C(x)$  and  $D(x)$  which appear in Eqs. (A.3), the covariant divergence of Eqs. (A.4) is calculated obtaining

$$\begin{aligned} f^{\mu} = & -\frac{\partial A}{\partial x^{\alpha}}\phi_{;\mu}^{\mu}\phi_{;\alpha}^{\alpha} - [A(x) + 2B(x)]\phi_{;\mu}^{\mu;\alpha}\phi_{;\alpha}^{\alpha} - A(x)\phi_{;\mu}^{\mu}\phi_{;\alpha}^{\alpha;\alpha} - \frac{\partial B}{\partial x^{\beta}}g^{\mu\beta}\phi_{;\mu}^{\mu}\phi_{;\alpha}^{\alpha} \\ & - \frac{\partial C}{\partial x^{\alpha}}\phi_{;\mu}^{\mu;\alpha} - C(x)\phi_{;\alpha;\mu}^{\mu;\alpha} - \frac{\partial D}{\partial x^{\beta}}g^{\mu\beta}\phi_{;\mu}^{\mu;\alpha} - D(x)\phi_{;\mu}^{\mu;\alpha;\alpha}. \end{aligned} \quad (\text{A.6})$$

where Eqs. (A.3) and (A.5) have been used.

These are the equations of the dynamics of the scalar field that determine the four functions  $A(x)$ ,  $B(x)$ ,  $C(x)$  and  $D(x)$  if the external forces  $f^{\mu}$  are known.

## Reference Frames

The ten Eqs. (A.4) of the gravitational field are not independent but are related by the four Bianchi identities which reduces Eqs. (A.4) to six independent equations. To determine an unambiguous metric, it is necessary to add four more equations to fix the gauge, that is, the reference frame.

In the presence of gravity, it is also convenient to define a reference frame at rest  $S'$  as described in section 2. Since Eqs. (3) are the same as in a Gravitational field in the presence of external forces [35] according to the Principle of Equivalence, then in any reference frame, for example, the reference frame  $S'$ , the equations of motion of the fluid will be

$$f'^{\mu} = T_M^{\prime\mu\alpha}_{;\alpha} = \frac{\partial T_M^{\prime\mu\alpha}}{\partial x'^{\alpha}} + \Gamma_{\alpha\beta}^{\prime\alpha}T_M^{\prime\mu\beta} + \Gamma_{\alpha\beta}^{\prime\mu}T_M^{\prime\alpha\beta}. \quad (\text{A.7})$$

Now, the Eqs. (2) must be also verified in the reference frame at rest  $S'$

$$f'^{\mu} = \frac{\partial T_M^{\prime\mu\alpha}}{\partial x'^{\alpha}}. \quad (\text{A.8})$$

This equates, in view of Eqs. (A.7) to choose a gauge such that

$$\Gamma'_{\alpha\beta}{}^{\mu} T'_{M\mu\beta} + \Gamma'_{\alpha\beta}{}^{\mu} T'_{M\alpha\beta} = 0. \quad (\text{A.9})$$

These four equations that determine the gauge, that is, the reference frame  $S'$ , are not covariant so they are not valid in any reference frame, but only in  $S'$ .

The moving reference frame  $S$  is chosen, as in section 2, such that the fluid, which originates the gravitational field, is at rest<sup>8</sup>

$$\frac{dx^i}{d\tau} = 0. \quad (\text{A.10})$$

In addition, to fix completely the reference frame  $S$ , it is necessary to add another condition that it can be obtained as in Eq. (9) by imposing that the time-time component of the metric must be the same in any reference frame

$$g_{tt} = g'_{tt} \quad (\text{A.11})$$

where  $g'_{tt}$  is the time-time component of the metric in the reference frame  $S'$  at rest.

The three Eqs. (A.10) together with Eq. (A.11) fix another gauge, that is, the reference frame  $S$ .

Once the reference frames have been fixed and the metric is known in  $S'$ , we can proceed as in the previous sections.

### Relevance of the Scalar Field

The introduction of the scalar field  $\phi$  gives the Minkowski metric Eqs. (1) as a solution of Einstein's Eqs. (A.4) in the gauge Eqs. (A.9). This solution is adequate in the absence of gravity. In practice, this happens when the gravitational field created by the source is so weak that it barely curves space-time. In this situation, in any reference frame, it is verified

$$R^{\mu\nu} = 0 \quad (\text{A.12})$$

and

$$R = g_{\alpha\beta} R^{\alpha\beta} = 0. \quad (\text{A.13})$$

It can be shown from Eqs. (A.12), (A.13) and (A.4)

$$T_M^{\mu\nu} = -T_{\phi}^{\mu\nu}, \quad (\text{A.14})$$

so the  $\phi$  field is especially important.

In vacuum, the energy-momentum tensor of the matter is zero

$$T_M^{\mu\nu} = 0 \quad (\text{A.15})$$

and Eq. (A.2) will be written as

$$\phi_{;\alpha}^{\alpha} = 0 \quad (\text{A.16})$$

while Eqs. (A.5) give

$$f^{\mu} = 0. \quad (\text{A.17})$$

Introducing Eqs. (A.16) and (A.17) into Eqs. (A.6) we obtain

---

<sup>8</sup> It is possible to find a reference frame like this because external forces can apply in such a way that all the particles of the fluid are at rest.

$$0 = \frac{\partial A}{\partial x^\alpha} \phi_{;\alpha}^\mu \phi_{;\alpha}^\alpha + [A(x) + 2B(x)] \phi_{;\alpha}^{\mu;\alpha} \phi_{;\alpha} + \frac{\partial B}{\partial x^\beta} g^{\mu\beta} \phi_{;\alpha}^\alpha \phi_{;\alpha} + \frac{\partial C}{\partial x^\alpha} \phi_{;\alpha}^{\mu;\alpha} + C(x) \phi_{;\alpha;\alpha}^{\alpha;\mu}. \quad (\text{A.18})$$

One solution of these equations, only valid on the outside of a static source, is

$$A(x) = B(x) = C(x) = 0. \quad (\text{A.19})$$

From Eqs. (A.16) and (A.19) the energy-momentum tensor (A.3) of the scalar field  $\phi$  in vacuum is

$$T_\phi^{\mu\nu} = 0. \quad (\text{A.20})$$

Equations (A.15) and (A.20) imply that in this case Eqs. (A.1) and (A.4) have the same solutions in vacuum and the scalar field does not alter the solutions of the Eqs. (A.1) that explain, among other phenomena, the dynamics of the planets. In other cases, their significance will have to be determined.

### Cosmological Dark Energy

Finally, it should be noted that in the absence of gravity or weak fields, the only way to produce acceleration over the fluid is, as it has been shown, by the application of external forces  $f^\mu$ . However, when the gravity generated by the fluid is strong, it can alone accelerate the fluid. Due to this, the energy-momentum tensor of the matter  $T_M^{\mu\nu}$  can not be conserved and the scalar field  $\phi$  can be present even if only gravitational forces act on the fluid. In this case, the introduction of the scalar field  $\phi$  into Einstein's Eqs. (A.4) through energy-momentum tensor Eqs. (A.3) adds four independent functions to the field equations. If the mass-energy of the matter is conserved

$$T_M^{t\alpha}_{;\alpha} = 0 \quad (\text{A.21})$$

only three degrees of freedom remain. This has important cosmological implications because three of the four functions  $A(x)$ ,  $B(x)$ ,  $C(x)$  and  $D(x)$ , of Eqs. (A.3), are independent and they can be adjusted to the observational data of accelerated expansion of the Universe.

## CONCLUSIONS

The exposed developments demonstrate the theoretical possibility of travelling to the past, in the case of a macroscopic perfect fluid without pressure in the absence of gravity and without violating the laws of Special Relativity. This is only possible if the fluid is accelerated to the speed of light. To demonstrate this, a relativistic treatment of rotation has been made using the principle of General Covariance, which has proved to be practical. Finally, it is necessary to introduce a Scalar Field in Einstein's equations to explain relativistic dynamics satisfactorily. This Scalar Field is the origin of the inertia and it has made possible to generalize the formalism to the case in which the fluid generates an appreciable Gravitational Field.

## ACKNOWLEDGEMENTS

This paper would not have been possible without the help of my friends Juan Miguel Reyes Montes and Manuel Saiz-Pardo Lizaso. Thank you for their collaboration and my gratitude for them.

## REFERENCES

- [1] Gödel, K.: An example of a new type of cosmological solution of Einstein's field equations of Gravitation. *Rev. Mod. Phys.* **21**, 447–450 (1949).
- [2] Tomimatsu, A., Sato, H.: New series of exact solutions for gravitational fields of spinning masses. *Prog. Theor. Phys.* **50**, 95-110 (1973).
- [3] Tipler, F.J.: Rotating cylinders and the possibility of global causality violation. *Phys. Rev. D* **9**, 2203-2206 (1974).
- [4] Deserr, S., Jackiw, R., 't Hooft, G.: Physical cosmic strings do not generate closed timelike curves. *Phys. Rev. Lett.* **68**, 267-269 (1992).
- [5] Gott, J.R.: Closed timelike curves produced by pairs of moving cosmic strings: Exact solutions. *Phys. Rev. Lett.* **66**, 1126-1129 (1991).
- [6] Ori, A.: A class of time-machine solutions with a compact vacuum core. *Phys. Rev. Lett.* **95**, 021101 (2005).
- [7] Ori, A.: Must time-machine construction violate the weak energy condition? *Phys. Rev. Lett.* **71**, 2517-2520 (1993).
- [8] Echeverria, F., Klinkhammer, G., Thorne, K.S.: Billiard balls in wormhole spacetimes with closed timelike curves: Classical theory. *Phys. Rev. D* **44**, 1077-1099 (1991).
- [9] Everett, A.E.: Warp drive and causality. *Phys. Rev. D* **53**, 7365-7368 (1996).
- [10] Everett, A.E., Roman, T.A.: Superluminal subway: The krasnikov tube. *Phys. Rev. D* **56**, 2100-2108 (1997).
- [11] Friedman, J., Morris, M.S., Novikov, I.D., Echeverria, F., Klinkhammer, G., Thorne, K.S., Yurtsever, U.: Cauchy problem in spacetimes with closed timelike curves. *Phys. Rev. D* **42**, 1915-1930 (1990).
- [12] Alcubierre, M.: The warp drive: Hyper-fast travel within general relativity. *Class. Quantum. Grav.* **11**, L73-L77 (1994).
- [13] Gonzalez-Diaz, P.F.: Warp drive space-time. *Phys. Rev. D* **62**, 044005 (2000).
- [14] Krasnikov, S.V.: Hyperfast travel in general relativity. *Phys. Rev. D* **57**, 4760-4766 (1998).
- [15] Lobo, F.S.N.: arXiv:1008.1127v1.
- [16] Morris, M.S., Thorne, K.S., Yurtsever, U.: Wormholes, time machines, and the weak energy condition. *Phys. Rev. Lett.* **61**, 1446-1449 (1988).
- [17] Misner, C.W., Thorne, K.S., Wheeler, J.A.: *Gravitation*, pp. 163-176. W. H. Freeman and Co., San Francisco (1973).
- [18] Ni, W.-T.: On the Proper reference frame and local coordinates of an accelerated observer in Special Relativity. *Chin.J.Phys.* **15**, 51-55 (1977).
- [19] Ni, W.-T, Zimmermann, M.: Inertial and gravitational effects in the proper reference frame of an accelerated, rotating observer. *Phys. Rev. D* **17**, 1473-1476 (1978).
- [20] Brumberg, V.A., Kopeikin, S.M.: Reference Frames, vol. 154 of *Astrophysics and Space Science Library*, pp. 115–141. J. Kovalevsky, I. I. Mueller, & B. Kolaczek (1989).



- [21] Damour, T., Soffel, M., Xu, C.: General relativistic celestial mechanics. 1. Method and definition of reference systems. *Phys. Rev. D* **43**, 3273-3307 (1991).
- [22] Damour, T., Soffel, M., Xu, C.: General relativistic celestial mechanics. 2. Translational equations of motion. *Phys. Rev. D* **45**, 1017-1044 (1992).
- [23] Damour, T., Soffel, M., Xu, C.: General relativistic celestial mechanics. 3. Rotational equations of motion. *Phys. Rev. D* **47**, 3124-3135 (1993).
- [24] Damour, T., Soffel, M., Xu, C.: General relativistic celestial mechanics. 4. Theory of satellite motion. *Phys. Rev. D* **49**, 618-635 (1994).
- [25] Kopeikin, S.M.: Celestial coordinate reference systems in curved space-time. *Celestial Mech.* **44**, 87-115 (1988).
- [26] Brumberg, V.A., Kopejkin, S.M.: Relativistic reference systems and motion of test bodies in the vicinity of the earth. *Nuovo Cim. B Serie* **103**, 63-98 (1989).
- [27] Kopeikin, S.M., Vlasov, I.: Parameterized post-Newtonian theory of reference frames, multipolar expansions and equations of motion in the N-body problem. *Phys. Rept.* **400**, 209-318 (2004).
- [28] Klauber, R.D.: Relativistic rotation: A comparison of theories. *Found. Phys.* **37**, 198-252 (2007).
- [29] Benedetto, E., Feleppa, F., Licata, I., Moradpour, H., Corda, C.: On the general relativistic framework of the Sagnac effect. *Eur. Phys. J. C* **79**, 187 (2019).
- [30] Corda, C.: New proof of general relativity through the correct physical interpretation of the Mössbauer rotor experiment. *Int.J.Mod.Phys. D* **27**, 1847016 (2018).
- [31] Nouri-Zonoz, M., Ramezani-Aval, H., Gharechahi, R.: On Franklin's relativistic rotational transformation and its modification. *Eur. Phys. J. C* **74**, 3098 (2014).
- [32] Soler, D.: Reference frames and rigid motions in relativity: Applications. *Found. Phys.* **36**, 1718-1735 (2006).
- [33] Klauber, R.D.: Analysis of the Anomalous Brilliet and Hall Experimental Result. *Found. Phys. Lett.* **17**, 125-147 (2004).
- [34] Kassner, K.: Spatial geometry of the rotating disk and its non-rotating counterpart. *Am. J. Phys.* **80**, 772-784 (2012).
- [35] Weinberg, S.: *Gravitation and Cosmology*. Wiley, New York (1972).
- [36] French, A.P.: *Special Relativity*, p. 71. Massachusetts Institute of Technology (1968).
- [37] Brans, C., Dicke, R.H.: Mach's Principle and a relativistic theory of Gravitation. *Phys. Rev.* **124**, 925-935 (1961).